

## MATH 147, SPRING 2021: FINAL EXAM PRACTICE PROBLEMS

Below are problems to practice for the final exam. The problems below, [together with the problems from the three midterm exams](#), are a good representation of what to expect on the final exam. There will also be a few short answer questions on the final exam.

1. Let  $f(x, y) = \begin{cases} x^2 + y^2, & \text{if } x^2 + y^2 < 1 \\ 1, & \text{if } x^2 + y^2 \geq 1. \end{cases}$  Determine at which points  $f(x, y)$  is continuous.

2. Show that the function  $f(x, y) = \begin{cases} \frac{2^x - 1}{xy} \sin(y), & \text{if } xy \neq 0 \\ \ln(2), & \text{if } xy = 0 \end{cases}$  is continuous at  $(0, 0)$ .

3. Use the limit definition to show that  $f(x, y) = 5x + 4y^2$  is differentiable at  $(2, 1)$ .

4. From class, we saw that if the first order partial derivatives of  $f(x, y)$  are continuous in a neighborhood of  $(a, b)$ , then  $f(x, y)$  is differentiable at  $(a, b)$ . This problem shows why those conditions are necessary. Let

$$f(x, y) = \begin{cases} \frac{2xy(x+y)}{x^2+y^2}, & \text{if } (x, y) \neq (0, 0) \\ 0, & \text{if } (x, y) = (0, 0) \end{cases}.$$

Show that:

- (i)  $f(x, y)$  is continuous at  $(0, 0)$ .
- (ii) Use the limit definitions to show that  $f_x(0, 0)$  and  $f_y(0, 0)$  exist and are equal to 0.
- (iii) Conclude that  $L(x, y) = 0$ .
- (iv) Show that  $f(x, y)$  is not differentiable at  $(0, 0)$ .
- (v) Show that  $f_x(x, y)$  is not continuous at  $(0, 0)$ .


5. Find and classify the critical points for  $f(x, y) = x^4 - 4xy + 2y^2$ .

6. Find the absolute maximum and absolute minimum values of  $f(x, y) = x^2y$  on the closed and bounded set  $D : 0 \leq 4x^2 + 9y^2 \leq 36$ .

7. Let  $S$  be the surface parametrized by  $G(u, v) = (2u \sin(\frac{v}{2}), 2u \cos(\frac{v}{2}), 3v)$ , with  $0 \leq u \leq 1$  and  $0 \leq v \leq 2\pi$ .

- (i) Find the tangent plane to  $S$  at the point  $P = G(1, \frac{\pi}{3})$ .
- (ii) Find the surface area of  $S$ .

8. Let  $C$  be a curve from the point  $P$  to the point  $Q$  in the  $xy$ -plane. Let  $\mathcal{R}$  be the region enclosed by  $C$  and the two radial lines from the origin to  $P$  and  $Q$ . (See the figure below.) Use Green's Theorem to show that  $\int_C \mathbf{F} \cdot d\mathbf{r}$  gives the area of  $\mathcal{R}$ , for  $\mathbf{F} = -\frac{y}{2}\mathbf{i} + \frac{x}{2}\mathbf{j}$ .




9. Let  $C$  be the triangle with vertices  $(1, 0, 0)$ ,  $(0, 2, 0)$ ,  $(0, 0, 1)$ . Compute  $\int_C \mathbf{F} \cdot d\mathbf{r}$ , for the vector field  $\mathbf{F} = (x^2 + yz, x + y, y - z^2)$ .

10. Let  $f(x, y) = \sqrt{|xy|}$ . Write out details showing:

- (a)  $\frac{\partial f}{\partial x}(0, 0)$  and  $\frac{\partial f}{\partial y}(0, 0)$  exist.

- (b)  $f(x, y)$  is not differentiable at  $(0, 0)$ .  
 (c) Part (b) does not contradict part (a).

11. Evaluate  $\int \int_S \text{Curl} \mathbf{F} \cdot d\mathbf{S}$ , for  $\mathbf{F} = (-y + z \sin(x), x, z^3)$  and  $S$  the surface defined by the equation  $x^2 + \frac{y^2}{4} + z^2 + z^4 x^2 = 1$ , with  $z \geq 0$ .
12. Verify the Divergence Theorem for  $\mathbf{F} = (-x^2, y^2, -z^2)$  and  $S$  rectangular box  $[0, 3] \times [-1, 2] \times [1, 2]$ .
13. Let  $\mathbf{F} = (z^2, x^2, -y^2)$ . Evaluate  $\int_C \mathbf{F} \cdot d\mathbf{r}$ , where  $C$  is the path traversing counterclockwise the square with sides of length  $s$  centered at  $(x_0, y_0, 0)$ . Then divide this number by the area of the square and take the limit as  $s \rightarrow 0$ . Compare this with  $(\text{Curl} \mathbf{F})(x_0, y_0, 0) \cdot \mathbf{k}$ .
14. Let  $C$  be the curve obtained by intersecting the cylinder  $x^2 + y^2 = 1$  with the plane  $x + y + z = 1$ , and  $\mathbf{F} = -y^3 \mathbf{i} + x^3 \mathbf{j} + -z^3 \mathbf{k}$ . Set up the line integral  $\int_C \mathbf{F} \cdot d\mathbf{r}$  as a single integral over an interval of the form  $[a, b]$ . Now evaluate this line integral by using Stoke's Theorem.
15. Verify Stoke's Theorem for  $\mathbf{F} = (z^2, -y^2, 0)$  and  $C$  the square of side 1 oriented as shown, lying in the  $xz$ -plane and  $S$  the open box with sides  $S_1, S_2, S_3, S_4, S_5$ . What happens, if instead, you take  $S$  to be the square enclosed by  $C$ ?



16. Calculate, without using Stoke's Theorem,  $\int \int_{S_1} \nabla \times \mathbf{F} \cdot d\mathbf{S}$ , for  $\mathbf{F} = (3y^2 + 2y)\mathbf{i} + 3z^2\mathbf{j} + 3x^2\mathbf{k}$  and  $S_1$  the inverted cone  $z = 1 - \sqrt{x^2 + y^2}$ , with vertex  $(0, 0, 1)$ , and  $z \geq 0$ . Then calculate directly  $\int \int_{S_2} \nabla \times \mathbf{F} \cdot d\mathbf{S}$ , for  $S_2$  the unit disk in the  $xy$ -plane. The answers you get should be the same. This shows the consequence of Stoke's Theorem, that surface integrals of the curl of a vector field over surfaces sharing the same boundary are independent of the surface.