

MATH 147, SPRING 2021: FINAL EXAM PRACTICE PROBLEMS

Below are problems to practice for the final exam. The problems below, [together with the problems from the three midterm exams](#), are a good representation of what to expect on the final exam. There will also be a few short answer questions on the final exam.

1. Let  $f(x, y) = \begin{cases} x^2 + y^2, & \text{if } x^2 + y^2 < 1 \\ 1, & \text{if } x^2 + y^2 \geq 1. \end{cases}$  Determine at which points  $f(x, y)$  is continuous.

**Solution.** Note that  $g(x, y) = x^2 + y^2$  and  $h(x, y) = 1$  are both continuous on all of  $\mathbb{R}^2$ . Thus, if we let  $D$  denote the unit disk  $0 \leq x^2 + y^2 \leq 1$ , then  $g(x, y)$  is continuous on the interior of  $D$  and  $h(x, y)$  is continuous on  $\mathbb{R}^2 \setminus D$ , and thus  $f(x, y)$  is continuous on both the interior of  $D$  and  $\mathbb{R}^2 \setminus D$ . For points  $(a, b)$  on the boundary of  $D$ ,  $a^2 + b^2 = 1$ , and we can consider  $\lim_{(x,y) \rightarrow (a,b)} f(x, y)$ . Let  $\epsilon > 0$ . Since  $g(x, y)$ , as a function on  $\mathbb{R}^2$ , is continuous at  $(a, b)$  there exists  $\delta > 0$  such that  $\|(x, y) - (a, b)\| < \delta$  implies  $|g(x, y) - g(a, b)| = |g(x, y) - 1| < \epsilon$ . Taking the same  $\delta$ , if  $\|(x, y) - (a, b)\| < \delta$  and  $(x, y) \in D$ , then  $g(x, y) = f(x, y)$ , which gives  $|f(x, y) - f(a, b)| = |g(x, y) - 1| < \epsilon$ . If  $(x, y) \notin D$ , then  $f(x, y) - f(a, b) = 1 - 1 = 0$ , so  $|f(x, y) - f(a, b)| < \epsilon$ . Thus  $f(x, y)$  is continuous at  $(a, b)$ .

2. Show that the function  $f(x, y) = \begin{cases} \frac{2^x - 1}{xy}(\sin(y)), & \text{if } xy \neq 0 \\ \ln(2), & \text{if } xy = 0 \end{cases}$  is continuous at  $(0, 0)$ .

**Solution.** Set  $g(x) = \begin{cases} \frac{2^x - 1}{x} & \text{if } x \neq 0 \\ \ln(2) & \text{if } x = 0 \end{cases}$  and  $h(y) = \begin{cases} \frac{\sin(y)}{y} & \text{if } y \neq 0 \\ 1 & \text{if } y = 0 \end{cases}$ . Then  $f(x, y) = g(x)h(y)$ .

Thus,  $\lim_{(x,y) \rightarrow (0,0)} f(x, y) = \{\lim_{x \rightarrow 0} g(x)\} \cdot \{\lim_{y \rightarrow 0} h(y)\}$ . By L'Hospital's Rule,  $\lim_{x \rightarrow 0} g(x) = \ln(2)$  and  $\lim_{y \rightarrow 0} h(y) = 1$ . Therefore,  $\lim_{(x,y) \rightarrow (0,0)} f(x, y) = \ln(2)$ .

3. Use the limit definition to show that  $f(x, y) = 5x + 4y^2$  is differentiable at  $(2, 1)$ .

**Solution.**  $\frac{\partial f}{\partial x}(2, 1) = 5$ ,  $\frac{\partial f}{\partial y}(2, 1) = 8$ , and  $f(2, 1) = 14$ , so  $L(x, y) = 5(x - 2) + 8(y - 1) + 14$ . Therefore,

$$\begin{aligned} f(x, y) - L(x, y) &= 5x + 4y^2 - (5x - 10 + 8y - 8 + 14) \\ &= 4y^2 - 8y + 4 \\ &= 4(y - 1)^2. \end{aligned}$$

Thus,  $\frac{f(x, y) - L(x, y)}{\sqrt{(x-2)^2 + (y-1)^2}} = \frac{4(y-1)^2}{\sqrt{(x-2)^2 + (y-1)^2}} \leq \frac{4(y-1)^2}{\sqrt{(y-1)^2}} = 4|y - 1|$ . Therefore,

$$\lim_{(x,y) \rightarrow (2,1)} \frac{f(x, y) - L(x, y)}{\sqrt{(x-2)^2 + (y-1)^2}} \leq \lim_{y \rightarrow 1} 4|y - 1| = 0,$$

which shows that  $f(x, y)$  is differentiable at  $(2, 1)$ .

4. From class, we saw that if the first order partial derivatives of  $f(x, y)$  are continuous in a neighborhood of  $(a, b)$ , then  $f(x, y)$  is differentiable at  $(a, b)$ . This problem shows why those conditions are necessary. Let

$$g(x, y) = \begin{cases} \frac{2xy(x+y)}{x^2+y^2}, & \text{if } (x, y) \neq (0, 0) \\ 0, & \text{if } (x, y) = (0, 0) \end{cases}.$$

Show that:

- (i)  $g(x, y)$  is continuous at  $(0, 0)$ .
- (ii) Use the limit definitions to show that  $g_x(0, 0)$  and  $g_y(0, 0)$  exist and are equal to 0.
- (iii) Conclude that  $L(x, y) = 0$ .
- (iv) Show that  $g(x, y)$  is not differentiable at  $(0, 0)$ .
- (v) Show that  $g_x(x, y)$  is not continuous at  $(0, 0)$ .

**Solution.** (i) Taking limits, we have.

$$\begin{aligned}\lim_{(x,y)\rightarrow(0,0)} g(x,y) &= \lim_{r\rightarrow 0} \frac{2r^2 \cos(\theta) \sin(\theta)(r \cos(\theta) + r \sin(\theta))}{r^2} \\ &= \lim_{r\rightarrow 0} r \cdot \{2 \cos(\theta) \sin(\theta)(\cos(\theta) + \sin(\theta))\} \\ &= 0 \\ &= g(0,0),\end{aligned}$$

so  $g(x,y)$  is continuous at  $(0,0)$ .

(ii)  $\frac{\partial g}{\partial x}(0,0) = \lim_{h\rightarrow 0} \frac{g(0+h,0)-g(0,0)}{h} = \lim_{h\rightarrow 0} \frac{0}{h} = 0$ .  $\frac{\partial g}{\partial y}(0,0) = \lim_{h\rightarrow 0} \frac{g(0,0+h)-g(0,0)}{h} = \lim_{h\rightarrow 0} \frac{0}{h} = 0$ .

(iii)  $L(x,y) = 0(x-0) + 0(y-0) = 0$ .

(iv) Thus,  $g(x,y) - L(x,y) = g(x,y)$ . If  $g(x,y)$  were differentiable at  $(0,0)$ , then the limit (when  $(x,y) \neq (0,0)$ )

$$\lim_{(x,y)\rightarrow(0,0)} \frac{g(x,y)}{\sqrt{x^2+y^2}} = \lim_{(x,y)\rightarrow(0,0)} \frac{2xy(x+y)}{(x^2+y^2)^{\frac{3}{2}}}$$

should equal zero. If we take the limit along the line  $y = x$ , then  $\frac{g(x,y)}{\sqrt{x^2+y^2}} = \frac{4x^3}{2\sqrt{2}x^3} = \sqrt{2}$ , so the limit along the line  $y = x$  as  $x \rightarrow 0$  is not 0. Thus,  $g(x,y)$  is not differentiable at  $(0,0)$ .

(v) Differentiating the non-zero part of  $g(x,y)$  gives  $g_x(x,y) = \frac{-2x^2y^2+4xy^3+2y^4}{(x^2+y^2)^2}$ . If we take the limit along the line  $y = 0$ , then  $\lim_{(x,y)\rightarrow(0,0)} g_x(x,y) = 0$ , while if we take the limit along the line  $x = 0$ , the limit becomes 2. Thus,  $\lim_{(x,y)\rightarrow(0,0)} g_x(x,y)$  does not exist, so that  $g_x(x,y)$  is not continuous at  $(0,0)$ .

5. Find and classify the critical points for  $f(x,y) = x^4 - 4xy + 2y^2$ .

**Solution.** To find critical points we solve

$$\begin{aligned}f_x &= 4x^3 - 4y = 0 \\ f_y &= -4x + 4y = 0.\end{aligned}$$

From the second equation, we get  $x = y$ . Using this in the first equation gives  $4x^3 - 4x = 0$ , so that  $x = 0, -1, 1$ . Thus, the critical points are  $(0,0)$ ,  $(-1,-1)$ , and  $(1,1)$ . For the discriminant, we have

$$D = f_{xx}f_{yy} - f_{xy}^2 = (12x^2)4 - (-4)^2 = 48x^2 - 16.$$

For  $(0,0)$ :  $D(0,0) = -16$ , so that  $f(x,y)$  has a saddle point at  $(0,0)$ .

For  $(-1,-1)$ :  $D(-1,-1) = 32 > 0$  and  $f_{xx}(-1,-1) = 12 > 0$ . Thus,  $f(x,y)$  has a relative minimum value (of -3) at  $(-1,-1)$ .

For  $(1,1)$ :  $D(1,1) = 32 > 0$  and  $f_{xx}(1,1) = 12 > 0$ . Thus,  $f(x,y)$  has a relative minimum value (of -3) at  $(1,1)$ .

6. Find the absolute maximum and absolute minimum values of  $f(x,y) = x^2y$  on the closed and bounded set  $D : 0 \leq 4x^2 + 9y^2 \leq 36$ .

**Solution.** Solving

$$\begin{aligned}f_x &= 2xy = 0 \\ f_y &= x^2 = 0\end{aligned}$$

we see that  $x = 0$ , and  $y$  can be any real number. Thus, critical points in the interior of  $D$  are of the form  $(0,y)$  with  $0 \leq 9y^2 \leq 36$ . However,  $f(0,y) = 0$  for all such points. On the boundary of  $D$ , we must

maximize and minimize  $f(x, y)$  subject to the constraint  $4x^2 + 9y^2 = 36$ . Calling this equation  $g(x, y)$ , we set  $\nabla f = \lambda \nabla g$  and solve the resulting system of equations

$$\begin{aligned} 2xy &= \lambda 8x \\ x^2 &= \lambda 18y \\ 4x^2 + 9y^2 &= 36. \end{aligned}$$

Notice that if  $x$  or  $y$  equal 0, then  $f(x, y) = 0$ . So we can assume neither  $x$  nor  $y$  is zero. Dividing the first equation by  $2x$  give  $y = 4\lambda$ . Using this in the second equation gives  $x^2 = 72\lambda^2$ . Putting both of these into the constraint equations gives  $4(72\lambda^2) + 9(4\lambda)^2 = 36$ , so that

$$3\lambda^2 = \frac{36}{144},$$

so that  $\lambda = \pm \frac{1}{\sqrt{12}}$ . Thus,  $y = \pm \frac{4}{\sqrt{12}}$  and  $x = \pm \sqrt{6}$ . Thus, substituting these into  $x^2y$  gives  $\pm \frac{24}{\sqrt{12}} = \pm 4\sqrt{3}$ . Thus, on the domain  $D$ , the maximum value of  $f(x, y)$  is  $4\sqrt{3}$  and the minimum value is  $-4\sqrt{3}$ . Note that the critical points  $(0, y)$  do not determine a minimum or maximum value of  $f(x, y)$  on  $D$ .

7. Let  $S$  be the surface parametrized by  $G(u, v) = (2u \sin(\frac{v}{2}), 2u \cos(\frac{v}{2}), 3v)$ , with  $0 \leq u \leq 1$  and  $0 \leq v \leq 2\pi$ .

- (i) Find the tangent plane to  $S$  at the point  $P = G(1, \frac{\pi}{3})$ .
- (ii) Find the surface area of  $S$ .

**Solution.**

$$\mathbf{T}_u \times \mathbf{T}_v = \begin{vmatrix} i & j & k \\ 2 \sin(v/2) & 2 \cos(v/2) & 0 \\ u \cos(v/2) & -u \sin(v/2) & 3 \end{vmatrix} = (6 \cos(v/2), -6 \sin(v/2), -2u).$$


Therefore,  $\mathbf{T}_u \times \mathbf{T}_v(1, \frac{\pi}{3}) = (3\sqrt{3}, -3, -2)$ , since  $G(1, \frac{\pi}{3}) = (1, \sqrt{3}, \pi)$ , for the tangent plane we have:

$$3\sqrt{3}(x - 1) - 3(y - \sqrt{3}) - 2(z - \pi) = 0.$$

For the surface area, we have  $\|\mathbf{T}_u \times \mathbf{T}_v\| = \sqrt{36 + 4u^2} = 2\sqrt{9 + u^2}$ .

$$\begin{aligned} \text{Surface area} &= \int \int_D \|\mathbf{T}_u \times \mathbf{T}_v\| \, dA \\ &= \int_0^{2\pi} \int_0^1 2\sqrt{9 + u^2} \, dudv \\ &= 4\pi \int_0^1 \sqrt{9 + u^2} \, du \\ &= 4\pi \left\{ \frac{u}{2} \sqrt{9 + u^2} + \frac{9}{2} \ln |u + \sqrt{9 + u^2}| \right\} \Big|_0^1 \quad (\text{using a table of integrals}) \\ &= 4\pi \left\{ \frac{\sqrt{10}}{2} + \frac{9}{2} \ln(1 + \sqrt{10}) - \frac{9}{2} \ln(3) \right\} \\ &= 4\pi \left\{ \frac{\sqrt{10}}{2} + \frac{9}{2} \ln\left(\frac{1 + \sqrt{10}}{3}\right) \right\}. \end{aligned}$$

8. Let  $C$  be a curve from the point  $P$  to the point  $Q$  in the  $xy$ -plane. Let  $\mathcal{R}$  be the region enclosed by  $C$  and the two radial lines from the origin to  $P$  and  $Q$ . (See the figure below.) Use Green's Theorem to show that  $\int_C \mathbf{F} \cdot d\mathbf{r}$  gives the area of  $\mathcal{R}$ , for  $\mathbf{F} = -\frac{y}{2}\mathbf{i} + \frac{x}{2}\mathbf{j}$ .



**Solution.** Let  $D$  denote the oriented closed curve forming the boundary of  $\mathbb{R}$ , namely, the line segment from the origin to  $P$ , followed by  $C$ , followed by the line segment from  $Q$  to the origin. We take  $\mathbf{F} = (\frac{-1}{2}y, \frac{1}{2}x)$ , so that by Green's Theorem,  $\int_D \mathbf{F} \cdot d\mathbf{r}$  equals the area enclosed by the closed curve  $D$ . Let  $C_1$  denote the line segment from  $(0,0)$  to  $P = (a,b)$ ,  $C_2$  denote the line segment from  $Q = (c,d)$  to  $(0,0)$ , and  $C$  the curve given in the illustration. Let  $D = C_1 \cup C \cup C_2$  so that

$$\text{area}(R) = \int_D \mathbf{F} \cdot d\mathbf{r} = \int_{C_1} \mathbf{F} \cdot d\mathbf{r} + \int_C \mathbf{F} \cdot d\mathbf{r} + \int_{C_2} \mathbf{F} \cdot d\mathbf{r}.$$

We have to show  $\int_{C_1} \mathbf{F} \cdot d\mathbf{r} = \int_{C_2} \mathbf{F} \cdot d\mathbf{r} = 0$ .

Take  $\mathbf{r}_1(t) = (at, bt)$ ,  $0 \leq t \leq 1$  for the parametrization of  $C_1$ . Then  $\mathbf{F}(\mathbf{r}(t)) = (-\frac{1}{2}bt, \frac{1}{2}at)$  and  $\mathbf{r}'(t) = (a, b)$ . Therefore  $\mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) = -\frac{1}{2}abt + \frac{1}{2}abt = 0$ . So  $\int_{C_1} \mathbf{F} \cdot d\mathbf{r} = 0$ . The calculation showing  $\int_{C_2} \mathbf{F} \cdot d\mathbf{r} = 0$  is similar.

9. Let  $C$  be the triangle with vertices  $(1,0,0)$ ,  $(0,2,0)$ ,  $(0,0,1)$ . Compute  $\int_C \mathbf{F} \cdot d\mathbf{r}$ , for the vector field  $\mathbf{F} = (x^2 + yz, x + y, y - z^2)$ .

**Solution.** We have to integrate along each side of the triangle.  $C_1 : \mathbf{r}(t) = (1-t, 2t, 0)$ , with  $0 \leq t \leq 1$ .  $\mathbf{r}'(t) = (-1, 2, 0)$ .  $\mathbf{F}(\mathbf{r}(t)) = ((1-t)^2, 1+t, 2t)$ .

$$\begin{aligned} \int_{C_1} \mathbf{F} \cdot d\mathbf{r} &= \int_0^1 \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt \\ &= \int_0^1 -(1-t)^2 + 2 + 2t dt \\ &= \int_0^1 -1 + 2t - t^2 + 2 + 2t dt \\ &= \int_0^1 1 + 4t - t^2 dt \\ &= 1 + 2 - \frac{1}{3} = \frac{8}{3}. \end{aligned}$$

$C_2 : \mathbf{r}(t) = (0, 2-2t, t)$ ,  $0 \leq t \leq 1$ ,  $\mathbf{r}'(t) = (0, -2, 1)$ ,  $\mathbf{F}(\mathbf{r}(t)) = (2t - 2t^2, 2 - 2t, 2 - 2t - t^2)$ .

$$\begin{aligned} \int_{C_2} \mathbf{F} \cdot d\mathbf{r} &= \int_0^1 \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt \\ &= \int_0^1 -4 + 4t + 2 - 2t - t^2 dt \\ &= \int_0^1 -2 + 2t - t^2 dt \\ &= -2 + 1 - \frac{1}{3} = -\frac{4}{3}. \end{aligned}$$

$C_3 : \mathbf{r}(t) = (t, 0, 1-t), \mathbf{r}'(t) = (1, 0, -1), \mathbf{F}(\mathbf{r}(t)) = (t^2, t, -(1-t)^2).$

$$\begin{aligned} \int_{C_2} \mathbf{F} \cdot d\mathbf{r} &= \int_0^1 \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt \\ &= \int_0^1 t^2 + (1-t)^2 dt \\ &= \left\{ \frac{t^3}{3} - \frac{(1-t)^3}{3} \right\} \Big|_0^1 \\ &= \frac{1}{3} - \left(-\frac{1}{3}\right) = \frac{2}{3}. \end{aligned}$$

Therefore,

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_{C_1} \mathbf{F} \cdot d\mathbf{r} + \int_C \mathbf{F} \cdot d\mathbf{r} + \int_{C_2} \mathbf{F} \cdot d\mathbf{r} = \frac{8}{3} - \frac{4}{3} + \frac{2}{3} = 2.$$

10. Let  $f(x, y) = \sqrt{|xy|}$ . Write out details showing:

- (a)  $\frac{\partial f}{\partial x}(0, 0)$  and  $\frac{\partial f}{\partial y}(0, 0)$  exist.
- (b)  $f(x, y)$  is not differentiable at  $(0, 0)$ .
- (c) Part (b) does not contradict part (a).

**Solution.**

2.(a)  $\frac{\partial f}{\partial x}(0, 0) = \lim_{h \rightarrow 0} \frac{f(0+h, 0) - f(0, 0)}{h} = \lim_{h \rightarrow 0} \frac{0}{h} = 0$ . The calculation for  $\frac{\partial f}{\partial y}(0, 0)$  is similar.

(b) From (a), the linear approximation to  $f(x, y)$  at  $(0, 0)$  is  $L(x, y) = 0$ . Therefore,  $f(x, y) - L(x, y) = f(x, y)$ . In order for  $f(x, y)$  to be differentiable at  $(0, 0)$ , the limit

$$\lim_{(x, y) \rightarrow (0, 0)} \frac{f(x, y) - L(x, y)}{\sqrt{x^2 + y^2}} = \lim_{(x, y) \rightarrow (0, 0)} \frac{f(x, y)}{\sqrt{x^2 + y^2}}$$

should be 0. Taking the limit along the line  $y = x$ , we have

$$\lim_{(x, y) \rightarrow (0, 0)} \frac{f(x, y)}{\sqrt{x^2 + y^2}} = \lim_{x \rightarrow 0} \frac{\sqrt{|x^2|}}{\sqrt{2}\sqrt{x^2}} = \frac{1}{\sqrt{2}} \neq 0.$$

Therefore,  $f(x, y)$  is not differentiable at  $(0, 0)$ .

(c) Part (b) does not contradict part (a) because the first order partial derivatives of  $f(x, y)$  are not continuous at  $(0, 0)$ . In fact, the partial derivatives of  $f(x, y)$  are not defined at all points  $(x, y)$  near  $(0, 0)$ , so we cannot evaluate the required limit to test continuity. To see this, let's try to calculate  $\frac{\partial f}{\partial x}$  along the line  $x = 0$ , with  $y \neq 0$ .

$$\frac{\partial f}{\partial x}(0, y) = \lim_{h \rightarrow 0} \frac{f(0+h, y) - f(0, y)}{h} = \lim_{h \rightarrow 0} \frac{|hy| - 0}{h} = |y| \lim_{h \rightarrow 0} \frac{|h|}{h},$$

which does not exist.

11. Evaluate  $\int \int_S \text{Curl} \mathbf{F} \cdot d\mathbf{S}$ , for  $\mathbf{F} = (-y + z \sin(x), x, z^3)$  and  $S$  the surface defined by the equation  $x^2 + \frac{y^2}{4} + z^2 + z^4 x^2 = 1$ , with  $z \geq 0$ .

**Solution.** We use Stoke's Theorem. The surface lies above the  $xy$ -plane, and intersects the  $xy$ -plane along the ellipse  $x^2 + \frac{y^2}{4} = 1$ . By Stokes Theorem,  $\int \int_S \text{Curl} \mathbf{F} \cdot d\mathbf{S} = \int_C \mathbf{F} \cdot d\mathbf{r}$ , for  $C : (\cos(t), 2 \sin(t), 0)$ , with  $0 \leq t \leq 2\pi$ . Note that the orientation of  $C$  is consistent with what is required by Stoke's Theorem.

$\mathbf{r}'(t) = (-\sin(t), 2\cos(t), 0)$  and  $\mathbf{F}(\mathbf{r}(t)) = (-2\sin(t), \cos(t), 0)$ . Therefore,

$$\begin{aligned}\iint_S \text{Curl } \mathbf{F} \cdot d\mathbf{S} &= \int_C \mathbf{F} \cdot d\mathbf{r} \\ &= \int_0^{2\pi} \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt \\ &= \int_0^{2\pi} 2 dt \\ &= 4\pi.\end{aligned}$$

12. Verify the Divergence Theorem for  $\mathbf{F} = (-x^2, y^2, -z^2)$  and  $S$  rectangular box  $[0, 3] \times [-1, 2] \times [1, 2]$ .

**Solution.**  $\text{Div } (\mathbf{F}) = -2x + 2y - 2z$ , therefore if  $B$  is the solid contained in the given rectangular box,

$$\begin{aligned}\iiint_B \text{Div } \mathbf{F} dV &= \int_1^2 \int_{-1}^2 \int_0^3 -2x + 2y - 2z dx dy dz \\ &= \int_1^2 \int_{-1}^2 (-x^2 + 2xy - 2xz) \Big|_{x=0}^{x=3} dy dz \\ &= \int_1^2 \int_{-1}^2 -9 + 6y - 6z dy dz \\ &= \int_0^1 (-9y + 3y^2 - 6yz) \Big|_{y=-1}^{y=2} dz \\ &= \int_1^2 -18 - 18z dz \\ &= (-18z - 9z^2) \Big|_{z=1}^{z=2} \\ &= (-36 - 36) - (-18 - 9) \\ &= -45.\end{aligned}$$

To calculate the surface integral, we must sum the integrals over each face of the given rectangular box.

**Front Face,  $S_1$ :**  $S_1$  is given by  $(3, y, z)$ , with  $-1 \leq y \leq 2$ ,  $1 \leq z \leq 2$  and  $\mathbf{n} = i$ . We will see the bounds on  $y$  and  $z$  are just used to calculate the area of the front face. The same will hold for the other five faces. So:  $\mathbf{F}$  on  $S_1$  is given by  $(-9, y^2, -z^2)$  and thus  $\mathbf{F} \cdot \mathbf{n} = -9$ . Therefore,

$$\begin{aligned}\iint_{S_1} \mathbf{F} \cdot d\mathbf{S} &= \iint_{S_1} \mathbf{F} \cdot \mathbf{n} dS \\ &= \iint_{S_1} -9 dS \\ &= -9 \cdot \text{area}(S_1) \\ &= -9 \cdot 3 = -27.\end{aligned}$$

**Back Face,  $S_2$ :**  $S_2$  is given by  $(0, y, z)$ , with  $-1 \leq y \leq 2$ ,  $1 \leq z \leq 2$  and  $\mathbf{n} = -i$ .  $\mathbf{F}$  on  $S_2$  is given by  $(0, y^2, -z^2)$  and thus  $\mathbf{F} \cdot \mathbf{n} = 0$ . Therefore,  $\iint_{S_2} \mathbf{F} \cdot d\mathbf{S} = \iint_{S_2} \mathbf{F} \cdot \mathbf{n} dS = 0$ .

**Left Face,  $S_3$ :**  $S_3$  is given by  $(x, -1, z)$ , with  $0 \leq x \leq 3$ ,  $1 \leq z \leq 2$  and  $\mathbf{n} = -j$ .  $\mathbf{F}$  on  $S_3$  is given by  $(-x^2, 1, -z^2)$  and thus  $\mathbf{F} \cdot \mathbf{n} = -1$ . Therefore,

$$\begin{aligned} \iint_{S_3} \mathbf{F} \cdot d\mathbf{S} &= \iint_{S_3} \mathbf{F} \cdot \mathbf{n} \, dS \\ &= \iint_{S_3} -1 \, dS \\ &= -1 \cdot \text{area}(S_3) \\ &= -1 \cdot 3 = -3. \end{aligned}$$

**Right Face,  $S_4$ :**  $S_4$  is given by  $(x, 2, z)$ , with  $0 \leq x \leq 3$ ,  $1 \leq z \leq 2$  and  $\mathbf{n} = j$ .  $\mathbf{F}$  on  $S_4$  is given by  $(-x^2, 4, -z^2)$  and thus  $\mathbf{F} \cdot \mathbf{n} = 4$ . Therefore,

$$\begin{aligned} \iint_{S_4} \mathbf{F} \cdot d\mathbf{S} &= \iint_{S_4} \mathbf{F} \cdot \mathbf{n} \, dS \\ &= \iint_{S_4} 4 \, dS \\ &= 4 \cdot \text{area}(S_4) \\ &= 4 \cdot 3 = 12. \end{aligned}$$

**Top Face,  $S_5$ :**  $S_5$  is given by  $(x, y, 2)$ , with  $0 \leq x \leq 3$ ,  $-1 \leq y \leq 2$  and  $\mathbf{n} = k$ .  $\mathbf{F}$  on  $S_5$  is given by  $(-x^2, y^2, -4)$  and thus  $\mathbf{F} \cdot \mathbf{n} = -4$ . Therefore,

$$\begin{aligned} \iint_{S_5} \mathbf{F} \cdot d\mathbf{S} &= \iint_{S_5} \mathbf{F} \cdot \mathbf{n} \, dS \\ &= \iint_{S_5} -4 \, dS \\ &= -4 \cdot \text{area}(S_5) \\ &= -4 \cdot 9 = -36. \end{aligned}$$

**Bottom Face,  $S_6$ :**  $S_6$  is given by  $(x, y, 1)$ , with  $0 \leq x \leq 3$ ,  $-1 \leq y \leq 2$  and  $\mathbf{n} = -k$ .  $\mathbf{F}$  on  $S_6$  is given by  $(-x^2, y^2, -1)$  and thus  $\mathbf{F} \cdot \mathbf{n} = 1$ . Therefore,

$$\begin{aligned} \iint_{S_6} \mathbf{F} \cdot d\mathbf{S} &= \iint_{S_6} \mathbf{F} \cdot \mathbf{n} \, dS \\ &= \iint_{S_6} 1 \, dS \\ &= 1 \cdot \text{area}(S_6) \\ &= 1 \cdot 9 = 9. \end{aligned}$$

Putting these all together we have  $\int \int_S \mathbf{F} \cdot d\mathbf{S} = -27 + 0 - 3 + 12 - 36 + 9 = -45$ .

13. Let  $\mathbf{F} = (z^2, x^2, -y^2)$ . Evaluate  $\int_C \mathbf{F} \cdot d\mathbf{r}$ , where  $C$  is the path traversing counterclockwise the square with sides of length  $s$  centered at  $(x_0, y_0, 0)$ . Then divide this number by the area of the square and take the limit as  $s \rightarrow 0$ . Compare this with  $(\text{Curl } \mathbf{F})(x_0, y_0, 0) \cdot k$ .

**Solution.** In this problem we are using the limit definition to calculate  $(\text{Curl } \mathbf{F})(x_0, y_0, 0) \cdot k$ . For this, we must compute a line integral of  $\mathbf{F}$  over each side of the square  $C$ .

**Bottom side,  $C_1$ :**  $C_1$  is given by  $\mathbf{r}(t) = (x_0 - \frac{s}{2}, y_0 - \frac{s}{2}, 0) + t(s, 0, 0)$ , with  $0 \leq t \leq 1$ .  $\mathbf{r}'(t) = (s, 0, 0)$ ,  $\mathbf{F}(\mathbf{r}(t)) = (0, (x_0 - \frac{s}{2})^2, -(y_0 - \frac{s}{2})^2)$ .

$$\begin{aligned}\int_{C_1} \mathbf{F} \cdot d\mathbf{r} &= \int_0^1 \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt \\ &= \int_0^1 0 dt \\ &= 0.\end{aligned}$$

**Top side,  $C_2$ :**  $C_2$  is given by  $\mathbf{r}(t) = (x_0 + \frac{s}{2}, y_0 + \frac{s}{2}, 0) + t(-s, 0, 0)$ , with  $0 \leq t \leq 1$ .  $\mathbf{r}'(t) = (-s, 0, 0)$ ,  $\mathbf{F}(\mathbf{r}(t)) = (0, (x_0 + \frac{s}{2})^2, -(y_0 + \frac{s}{2})^2)$ .

$$\begin{aligned}\int_{C_2} \mathbf{F} \cdot d\mathbf{r} &= \int_0^1 \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt \\ &= \int_0^1 0 dt \\ &= 0.\end{aligned}$$

**Right side,  $C_3$ :**  $C_3$  is given by  $\mathbf{r}(t) = (x_0 + \frac{s}{2}, y_0 - \frac{s}{2}, 0) + t(0, s, 0)$ , with  $0 \leq t \leq 1$ .  $\mathbf{r}'(t) = (0, s, 0)$ ,  $\mathbf{F}(\mathbf{r}(t)) = (0, (x_0 + \frac{s}{2})^2, -(y_0 - \frac{s}{2})^2)$ .

$$\begin{aligned}\int_{C_3} \mathbf{F} \cdot d\mathbf{r} &= \int_0^1 \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt \\ &= \int_0^1 (x_0 + \frac{s}{2})^2 s dt \\ &= x_0^2 s + x_0 s^2 + \frac{s^3}{4}.\end{aligned}$$

**Left side,  $C_4$ :**  $C_4$  is given by  $\mathbf{r}(t) = (x_0 - \frac{s}{2}, y_0 + \frac{s}{2}, 0) + t(0, -s, 0)$ , with  $0 \leq t \leq 1$ .  $\mathbf{r}'(t) = (0, -s, 0)$ ,  $\mathbf{F}(\mathbf{r}(t)) = (0, (x_0 - \frac{s}{2})^2, -(y_0 + \frac{s}{2})^2)$ .

$$\begin{aligned}\int_{C_4} \mathbf{F} \cdot d\mathbf{r} &= \int_0^1 \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt \\ &= \int_0^1 -(x_0 - \frac{s}{2})^2 s dt \\ &= -x_0^2 s + x_0 s^2 - \frac{s^3}{4}.\end{aligned}$$

We now have

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_{C_1} \mathbf{F} \cdot d\mathbf{r} + \int_{C_2} \mathbf{F} \cdot d\mathbf{r} + \int_{C_3} \mathbf{F} \cdot d\mathbf{r} + \int_{C_4} \mathbf{F} \cdot d\mathbf{r} = 0 + 0 + (x_0^2 s + x_0 s^2 + \frac{s^3}{4}) + (-x_0^2 s + x_0 s^2 - \frac{s^3}{4}) = 2s^2 x_0.$$

Therefore,

$$\lim_{s \rightarrow 0} \frac{1}{\text{area}(S)} \int_C \mathbf{F} \cdot d\mathbf{r} = \lim_{s \rightarrow 0} \frac{1}{s^2} 2s^2 x_0 = 2x_0.$$

To Check:

$$\text{Curl } \mathbf{F} = \begin{vmatrix} i & j & k \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ z^2 & x^2 & -y^2 \end{vmatrix} = (-2y, 2z, 2x).$$

Thus,  $\text{Curl } \mathbf{F}(x_0, y_0, z_0) = (-2y_0, 2z_0, 2x_0)$  and  $\text{Curl } \mathbf{F}(x_0, y_0, z_0) \cdot \mathbf{k} = 2x_0$ .

14. Let  $C$  be the curve obtained by intersecting the cylinder  $x^2 + y^2 = 1$  with the plane  $x + y + z = 1$ , and  $\mathbf{F} = -y^3 \vec{i} + x^3 \vec{j} + -z^3 \vec{k}$ . Set up the line integral  $\int_C \mathbf{F} \cdot d\mathbf{r}$  as a single integral over an interval of the form  $[a, b]$ . Now evaluate this line integral by using Stoke's Theorem.



**Solution.** The curve  $C$  lies on the plane  $z = 1 - x - y$  but also on the cylinder  $x^2 + y^2 = 1$ . So a parametrization and tangent of  $C$  are

$$\mathbf{r}(t) = (\cos(t), \sin(t), 1 - \cos(t) - \sin(t)) \text{ and } \mathbf{r}'(t) = (-\sin(t), \cos(t), \sin(t) - \cos(t)).$$

$$\mathbf{F}(\mathbf{r}(t)) = (-\sin^3(t), \cos^3(t), -(1 - \cos(t) - \sin(t))^3).$$

$$\mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) = \sin^4(t) + \cos^4(t) - (1 - \cos(t) - \sin(t))^3 \cdot (\sin(t) - \cos(t)).$$

Integrating this last expression from 0 to  $2\pi$  is doable .... but not much fun.

To apply Stoke's Theorem we will integrate  $\nabla \times \mathbf{F}$  over  $S$ , that portion of the given plane lying above the disk  $D : 0 \leq x^2 + y^2 \leq 1$  in the  $xy$ -plane.

$$\nabla \times \mathbf{F} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ -y^3 & x^3 & -z^3 \end{vmatrix} = (3x^2 + 3y^2)\vec{k} = (0, 0, 3x^2 + 3y^2).$$

$S$  is given by  $G(u, v) = (u, v, 1 - u - v)$ , with  $0 \leq u^2 + v^2 \leq 1$ . Thus,

$$\mathbf{T}_u \times \mathbf{T}_v = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & 0 & -1 \\ 0 & 1 & -1 \end{vmatrix} = \vec{i} + \vec{j} + \vec{k} = (1, 1, 1).$$


Moreover,  $\nabla \times \mathbf{F}$  on  $S$  is  $(0, 0, 3u^2 + 3v^2)$ . Thus, on  $S$ ,

$$(\nabla \times \mathbf{F}) \cdot \mathbf{T}_u \times \mathbf{T}_v = (0, 0, 3u^2 + 3v^2) \cdot (1, 1, 1) = 3u^2 + 3v^2.$$

Therefore:

$$\begin{aligned} \int \int_S \nabla \times \mathbf{F} \cdot d\mathbf{S} &= \int \int_D 3u^2 + 3v^2 \, dA \\ &= 3 \int_0^{2\pi} \int_0^1 r^2 \cdot r \, dr d\theta \\ &= 6\pi \int_0^1 r^3 \, dr \\ &= \frac{6\pi}{4} = \frac{3\pi}{2}. \end{aligned}$$

15. Verify Stoke's Theorem for  $\mathbf{F} = (z^2, -y^2, 0)$  and  $C$  the square of side 1 oriented as shown, lying in the  $xz$ -plane and  $S$  the open box with sides  $S_1, S_2, S_3, S_4, S_5$ . What happens, if instead, you take  $S$  to be the square enclosed by  $C$ ?



**Solution.** Both terms in Stoke's Theorem require computing several integrals. We start by computing  $\int_C \mathbf{F} \cdot d\mathbf{r}$ , where  $C = C_1 \cup C_2 \cup C_3 \cup C_4$  is the curve indicated in the diagram.

$$C_1 : \mathbf{r}(t) = (1 - t, 0, 0), 0 \leq t \leq 1, \mathbf{r}'(t) = (-1, 0, 0), \mathbf{F}(\mathbf{r}(t)) = (0, 0, 0).$$

$$\begin{aligned} \int_{C_1} \mathbf{F} \cdot d\mathbf{r} &= \int_0^1 \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt \\ &= \int_0^1 0 dt \\ &= 0. \end{aligned}$$

$$C_2 : \mathbf{r}(t) = (0, 0, t), 0 \leq t \leq 1, \mathbf{r}'(t) = (0, 0, 1), \mathbf{F}(\mathbf{r}(t)) = (t^2, 0, 0).$$

$$\begin{aligned} \int_{C_2} \mathbf{F} \cdot d\mathbf{r} &= \int_0^1 \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt \\ &= \int_0^1 0 dt \\ &= 0. \end{aligned}$$

$$C_3 : \mathbf{r}(t) = (t, 0, 1), 0 \leq t \leq 1, \mathbf{r}'(t) = (1, 0, 0), \mathbf{F}(\mathbf{r}(t)) = (1, 0, 0).$$

$$\begin{aligned} \int_{C_3} \mathbf{F} \cdot d\mathbf{r} &= \int_0^1 \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt \\ &= \int_0^1 1 dt \\ &= 1. \end{aligned}$$

$$C_4 : \mathbf{r}(t) = (1, 0, 1 - t), 0 \leq t \leq 1, \mathbf{r}'(t) = (0, 0, -1), \mathbf{F}(\mathbf{r}(t)) = ((1 - t)^2, 0, 0).$$

$$\begin{aligned} \int_{C_4} \mathbf{F} \cdot d\mathbf{r} &= \int_0^1 \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt \\ &= \int_0^1 0 dt \\ &= 0. \end{aligned}$$

We now have,

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_{C_1} \mathbf{F} \cdot d\mathbf{r} + \int_{C_2} \mathbf{F} \cdot d\mathbf{r} + \int_{C_3} \mathbf{F} \cdot d\mathbf{r} + \int_{C_4} \mathbf{F} \cdot d\mathbf{r} = 0 + 0 + 1 + 0 = 1.$$

To calculate the curl of  $\mathbf{F}$

$$\text{Curl } \mathbf{F} = \begin{vmatrix} i & j & k \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ z^2 & -y^2 & 0 \end{vmatrix} = (0, 2z, 0).$$

For the surface integral  $\int \int_S \text{Curl } \mathbf{F} \cdot d\mathbf{S}$  is the sum of the surface integrals of the curl of  $\mathbf{F}$  over the five faces indicated in the diagram.

**Front Face,  $S_1$ :**  $S_1$  is given by  $(1, y, z)$ , with  $0 \leq y \leq 1$ ,  $0 \leq z \leq 1$  and  $\mathbf{n} = i$ . Curl  $\mathbf{F}$  on  $S_1$  is given by  $(0, 2z, 0)$  and thus  $\mathbf{F} \cdot \mathbf{n} = 0$ . Therefore,

$$\begin{aligned} \int \int_{S_1} \text{Curl } \mathbf{F} \cdot d\mathbf{S} &= \int \int_{S_1} \text{Curl } \mathbf{F} \cdot \mathbf{n} dS \\ &= \int \int_{S_1} 0 dS \\ &= 0. \end{aligned}$$

**Back Face,  $S_2$ :**  $S_2$  is given by  $(0, y, z)$ , with  $0 \leq y \leq 1$ ,  $0 \leq z \leq 1$  and  $\mathbf{n} = -i$ .  $\text{Curl } \mathbf{F}$  on  $S_2$  is given by  $(0, 2z, 0)$  and thus  $\mathbf{F} \cdot \mathbf{n} = 0$ . Therefore,

$$\begin{aligned} \iint_{S_2} \text{Curl } \mathbf{F} \cdot d\mathbf{S} &= \iint_{S_2} \text{Curl } \mathbf{F} \cdot \mathbf{n} \, dS \\ &= \iint_{S_3} 0 \, dS \\ &= 0. \end{aligned}$$

**Top Face,  $S_3$ :**  $S_3$  is given by  $(x, y, 1)$ , with  $0 \leq x \leq 1$ ,  $0 \leq y \leq 1$  and  $\mathbf{n} = k$ .  $\text{Curl } \mathbf{F}$  on  $S_3$  is given by  $(0, 2, 0)$  and thus  $\mathbf{F} \cdot \mathbf{n} = 0$ . Therefore,

$$\begin{aligned} \iint_{S_3} \text{Curl } \mathbf{F} \cdot d\mathbf{S} &= \iint_{S_3} \text{Curl } \mathbf{F} \cdot \mathbf{n} \, dS \\ &= \iint_{S_3} 0 \, dS \\ &= 0. \end{aligned}$$

**Bottom Face,  $S_4$ :**  $S_4$  is given by  $(x, y, 0)$ , with  $0 \leq x \leq 1$ ,  $0 \leq y \leq 1$  and  $\mathbf{n} = -k$ .  $\text{Curl } \mathbf{F}$  on  $S_4$  is given by  $(0, 0, 0)$  and thus  $\mathbf{F} \cdot \mathbf{n} = 0$ . Therefore,

$$\begin{aligned} \iint_{S_4} \text{Curl } \mathbf{F} \cdot d\mathbf{S} &= \iint_{S_4} \text{Curl } \mathbf{F} \cdot \mathbf{n} \, dS \\ &= \iint_{S_4} 0 \, dS \\ &= 0. \end{aligned}$$

**Right Face,  $S_5$ :**  $S_5$  is given by  $(x, 1, z)$ , with  $0 \leq x \leq 1$ ,  $0 \leq z \leq 1$  and  $\mathbf{n} = j$ .  $\text{Curl } \mathbf{F}$  on  $S_5$  is given by  $(0, 2z, 0)$  and thus  $\mathbf{F} \cdot \mathbf{n} = 2z$ . Therefore,

$$\begin{aligned} \iint_{S_5} \text{Curl } \mathbf{F} \cdot d\mathbf{S} &= \iint_{S_5} \text{Curl } \mathbf{F} \cdot \mathbf{n} \, dS \\ &= \iint_{S_5} 2z \, dS \\ &= \int_0^1 \int_0^1 2z \, dz \, dx \\ &= \int_0^1 2z \, dz \\ &= 1. \end{aligned}$$

Putting these all together we have  $\int \int_S \text{Curl } \mathbf{F} \cdot d\mathbf{S} = 0 + 0 + 0 + 0 + 1 = 1$ , as expected, thereby confirming Stoke's Theorem.

Finally, an important consequence of Stoke's theorem is that the surface integrals of  $\text{Curl } \mathbf{F}$  over two surfaces sharing a common oriented boundary are the same. Let  $S_0$  be the left face of the square in the diagram, so that  $S$  and  $S_0$  share the same **oriented** boundary.

$S_0$  is given by  $(x, 0, z)$ , with  $0 \leq x \leq 1$ ,  $0 \leq z \leq 1$  and  $\mathbf{n} = j$ . Note that  $j$  is the correct normal when considering  $S_0$  as an open, oriented surface with boundary  $C$ . If we were considering  $S_0$  as the 6th side of the cube, we would take  $-j$  as the unit normal.  $\text{Curl } \mathbf{F}$  on  $S_0$  is given by  $(0, 2z, 0)$  and thus  $\text{Curl } \mathbf{F} \cdot \mathbf{n} = 2z$ .

Therefore,

$$\begin{aligned}
 \iint_{S_0} \text{Curl } \mathbf{F} \cdot d\mathbf{S} &= \iint_{S_0} \mathbf{F} \cdot \mathbf{n} \, dS \\
 &= \iint_{S_0} 2z \, dS \\
 &= \int_0^1 \int_0^1 2z \, dz \, dx \\
 &= \int_0^1 2z \, dz \\
 &= 1 \\
 &= \iint_S \text{Curl } \mathbf{F} \cdot d\mathbf{S}.
 \end{aligned}$$

16. Calculate, without using Stoke's Theorem,  $\int \int_{S_1} \nabla \times \mathbf{F} \cdot d\mathbf{S}$ , for  $\mathbf{F} = (3y^2 + 2y)\vec{i} + 3z^2\vec{j} + 3x^2\vec{k}$  and  $S_1$  the inverted cone  $z = 1 - \sqrt{x^2 + y^2}$ , with vertex  $(0, 0, 1)$ , and  $z \geq 0$ . Then calculate directly  $\int \int_{S_2} \nabla \times \mathbf{F} \cdot d\mathbf{S}$ , for  $S_2$  the unit disk in the  $xy$ -plane. The answers you get should be the same. This shows the consequence of Stoke's Theorem, that surfaces integrals of the curl of a vector field over surfaces sharing the same boundary are independent of the surface.

**Solution.**  $\nabla \times \mathbf{F} = \begin{vmatrix} i & j & k \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 3y^2 + 2y & 3z^2 & 3x^2 \end{vmatrix} = (-6z, -6x, -6y - 2).$

If we integrate directly over the cone, we use the parametrization

$$G(u, v) = (v \cos(u), v \sin(u), 1 - v),$$

with  $0 \leq u \leq 2\pi$ ,  $0 \leq v \leq 1$ . It follows that

$$\nabla \times \mathbf{F}(G(u, v)) = (-6(1 - v), -6v \cos(u), -6v \sin(u) - 2).$$

Moreover,

$$\mathbf{T}_u \times \mathbf{T}_v = (v \cos(u), v \sin(u), -v).$$

Note that this vector, has a negative  $z$ -component, and thus is *inside* of the inverted cone. If we flatten the cone, this would point downward, and contradict the right hand thumb rule. Thus, we need to use the vector  $-(\mathbf{T}_u \times \mathbf{T}_v)$  when integrating  $\nabla \mathbf{F}$ . (An important point: We could also parametrize the inverted cone using  $H(u, v) = (u, v, 1 - \sqrt{u^2 + v^2})$ , with  $0 \leq u^2 + v^2 \leq 1$ , and in this case  $\mathbf{T}_u \times \mathbf{T}_v$  gives the correct normal vector.)

We now have

$$\{\nabla \times \mathbf{F}(G(u, v))\} \cdot -(\mathbf{T}_u \times \mathbf{T}_v) = 6v(1 - v) \cos(u) + 6v^2 \sin(u) \cos(u) - 6v^2 \sin(u) - 2v.$$

When we calculate  $\int \int_S (\nabla \times \mathbf{F}) \cdot d\mathbf{S}$ , first integrating with respect to  $u$ , the trig terms integrated from 0 to  $2\pi$  all become 0. We are left with,

$$\begin{aligned}
 \int_0^1 \int_0^{2\pi} -2v \, dudv &= 2\pi \int_0^1 -2v \, dv \\
 &= -4\pi \cdot \frac{v^2}{2} \Big|_0^1 \\
 &= -2\pi.
 \end{aligned}$$

To integrate over the disk we have  $G(u, v) = (u, v, 0)$  with  $0 \leq u^2 + v^2 \leq 1$  and  $\mathbf{T}_u \times \mathbf{T}_v = (0, 0, 1)$ .  $\nabla \times \mathbf{F}(G(u, v)) = (0, -6u, -6v - 2)$ , so  $\nabla \times \mathbf{F}(G(u, v)) \cdot \mathbf{T}_u \times \mathbf{T}_v = -6v - 2$ . Thus,

$$\begin{aligned}\iint_{S_2} (\nabla \times \mathbf{F}) \cdot d\mathbf{S} &= \iint_{S_2} -6v - 2 \, dA \\ &= \int_0^{2\pi} \int_0^1 (-6r \sin(\theta) - 2) \, r \, dr \, d\theta \\ &= \int_0^{2\pi} \int_0^1 -6r^2 \sin(\theta) - 2r \, dr \, d\theta \\ &= \int_0^{2\pi} -2 \sin(\theta) - 1 \, d\theta \\ &= -2\pi,\end{aligned}$$

which is what we want.